arXiv:1404.0458v1 [math.DG] 2 Apr 2014

REALIZATION OF AN EQUIVARIANT HOLOMORPHIC HERMITIAN LINE BUNDLE AS A QUILLEN DETERMINANT BUNDLE

INDRANIL BISWAS

Abstract. Let M be an irreducible smooth complex projective variety equipped with an action of a compact Lie group G, and let $(\mathcal{L}\,, h)$ be a G–equivariant holomorphic Hermitian line bundle on M. Given a compact connected Riemann surface X, we construct a G–equivariant holomorphic Hermitian line bundle $(L\,, H)$ on $X \times M$ (the action of G on X is trivial), such that the corresponding Quillen determinant line bundle $(\mathcal{Q}\,, h_Q)$, which is a G–equivariant holomorphic Hermitian line bundle on M, is isomorphic to the given G–equivariant holomorphic Hermitian line bundle $(\mathcal{L}\,, h)$. This proves a conjecture in [DM].

1. Introduction

This work was inspired by [DM] where the following result is proved. Let M be an irreducible smooth complex projective variety and $\mathcal{L}$ an ample line bundle on M equipped with a Hermitian structure h of positive curvature. There is a natural family of Cauchy–Riemann operators on $\mathbb{CP}^1$, parametrized by M, such that the corresponding Quillen determinant line bundle, which is a holomorphic Hermitian line bundle on M, is holomorphically isomorphic to a positive tensor power of $(\mathcal{L}\,, h)$ [DM, p. 785, Theorem 1.1]. It is conjectured in [DM] that an equivariant version also holds (see [DM, p. 793, § 5]).

Let M be as before. Assume that it is equipped with a C^∞ action of a compact Lie group G via holomorphic automorphisms of M. Let $(\mathcal{L}\,, h)$ be any G–equivariant holomorphic Hermitian line bundle on M.

Let X be a compact connected Riemann surface equipped with a Kähler form ω_X. Let L_0 be a holomorphic line bundle on X of degree $\text{genus}(X) - 2$ such that $H^0(X,\, L_0) = 0$. Fix a Hermitian structure h_0 on L_0.

The action of G on M and the trivial action of G on X together produce an action of G on $X \times M$. Let p_1 and p_2 be the projections of $X \times M$ on X and M respectively. Consider the Hermitian structure $H := (p_1^* h_0) \otimes (p_2^* h)$ on the holomorphic line bundle

$$L := (p_1^* L_0) \otimes (p_2^* \mathcal{L})$$

over $X \times M$. The action of G on $\mathcal{L}$ and the trivial action of G on L_0 together produce an action of G on L, thus making L a G–equivariant holomorphic Hermitian line bundle on $X \times M$. We will consider $(L\,, H)$ as a family of holomorphic Hermitian line bundles o X.

2000 *Mathematics Subject Classification.* 58J52, 14H60.

Key words and phrases. Quillen determinant, equivariant bundle, fiber integral.

Let $(\mathcal{Q}\,, h_Q)$ be the Quillen determinant line bundle associated to the triple $(L\,,\, H\,, \omega_X)$. It is a G–equivariant holomorphic Hermitian line bundle on M.

We prove the following (see Theorem 2.4):

The two G–equivariant holomorphic Hermitian line bundle on M, namely $(\mathcal{L}\,, h)$ and $(\mathcal{Q}\,, h_Q)$, are isomorphic.

We note that this proves the earlier mentioned conjecture in [DM].

2. A holomorphic family and its determinant bundle

Let X be a compact connected Riemann surface. Let g be the genus of X. Fix a holomorphic line bundle L_0 on X of degree $g-2$ such that

$$H^0(X,\, L_0) \,=\, 0\,. \tag{2.1}$$

We note that such a line bundle exists. Indeed, if $g \,\leq\, 1$, then any holomorphic line bundle of degree $g-2$ works; if $g \,=\, 2$, then any nontrivial holomorphic line bundle of degree zero works; if $g \,\geq\, 2$, then any point outside the image of the Abel-Jacobi map $\text{Sym}^{g-2}(X) \,\longrightarrow\, \text{Pic}^{g-2}(X)$ works. From Riemann–Roch it follows that

$$\dim H^1(X,\, L_0) \,=\, 1\,. \tag{2.2}$$

Fix a C^∞ Hermitian structure h_0 on L_0. Also, fix a Kähler form ω_X on X.

Let M be a connected complex projective manifold (meaning a connected smooth complex projective variety). Assume that a compact Lie group G acts smoothly on M via holomorphic automorphisms. Let $(\mathcal{L}\,, h)$ be a G–equivariant holomorphic Hermitian line bundle on M. This means that the holomorphic line bundle L is equipped with an action of G such that

(1) for each element $z \,\in\, G$, the action of z on $\mathcal{L}$ is a holomorphic automorphism of the line bundle $\mathcal{L}$ over the automorphism of M given by the action of z on M,
(2) the action of G on $\mathcal{L}$ is C^∞ and it preserves h.

Let p_1 and p_2 be the projections on $X \times M$ to X and M respectively. Consider the holomorphic line bundle

$$L \,:=\, (p_1^* L_0) \otimes (p_2^* \mathcal{L}) \,\longrightarrow\, X \times M\,.$$

It is equipped with the Hermitian structure

$$H \,:=\, (p_1^* h_0) \otimes (p_2^* h)\,. \tag{2.3}$$

The action of G on M and the trivial action of G on X together define an action of G on $X \times M$. Similarly, the action of G on $\mathcal{L}$ and the trivial action of G on L_0 together define an action of G on L. This action of G on L clearly preserves H.

Consider $(L\,, H)$ as a family of holomorphic Hermitian line bundles o X parametrized by M. Let

$$(\mathcal{Q}\,, h_Q) \,\longrightarrow\, M$$

be the Quillen determinant line bundle associated to $(L\,,\,H\,,\omega_X)$ [Qu]. For any point $y\,\in\,M$, let $(L^y\,,H^y)$ be the holomorphic Hermitian line bundle on X obtained by restricting $(L\,,H)$ to $X\,\times\,\{y\}$. Note that $(L^y\,,H^y)$ is isomorphic to the holomorphic Hermitian line bundle $(L_0\,,h_0)$. We recall that the fiber $\mathcal{Q}_y$ is identified with the complex line $\bigwedge^{\rm top} H^0(X\,,L^y)^*\otimes\bigwedge^{\rm top} H^1(X,\,L^y)$ [Qu]. In view of (2.1) and (2.2), the fiber $\mathcal{Q}_y$ is identified with $H^1(X,\,L^y)$.

The action of G on L produces an action of G on $\mathcal{Q}$. The action of any $z\,\in\,G$ on $\mathcal{Q}$ is a holomorphic automorphism of the line bundle $\mathcal{Q}$ over the automorphism of M given by z. The action of G on $\mathcal{Q}$ preserves the Hermitian structure h_Q on $\mathcal{Q}$ because the action of G on L preserves H and the trivial action on X preserves ω_X.

Let

$$\xi\,:=\,M\times H^1(X,\,L_0)\,\longrightarrow\,M \tag{2.4}$$

be the holomorphically trivial line bundle with fiber $H^1(X,\,L_0)$ (see (2.2)). The trivial action of G on $H^1(X,\,L_0)$ and the action of G on M together define an action of G on ξ. The actions of G on $\mathcal{L}$ and ξ together produce an action of G on $\mathcal{L}\otimes\xi$ that is a lift of the action of G on M.

Lemma 2.1. *The holomorphic line bundle $\mathcal{Q}$ over M is identified with $\mathcal{L}\otimes\xi$. This identification is G–equivariant.*

Proof. From (2.1) it follows that

$$R^0p_{2*}L\,=\,0$$

(recall that L^y is isomorphic to L_0). By the projection formula [Ha, p. 253, Ex. 8.3], we have

$$R^1p_{2*}L\,=\,\mathcal{L}\otimes R^1p_{2*}(p_1^*L_0)\,.$$

But $R^1p_{2*}(p_1^*L_0)\,=\,\xi$. Therefore, we get an isomorphism

$$\tau\,:\,\mathcal{Q}\,:=\,{\rm Det}(L)\,=\,R^1p_{2*}L\,\stackrel{\sim}{\longrightarrow}\,\mathcal{L}\otimes\xi\,. \tag{2.5}$$

From the construction of τ it follows immediately that it intertwines the actions of G on $\mathcal{Q}$ and $\mathcal{L}\otimes\xi$. □

Let ∇^Q be the Chern connection on $\mathcal{Q}$ for the Hermitian structure h_Q. The curvature of ∇^Q will be denoted by $\mathcal{K}(\nabla^Q)$. The curvature $\mathcal{K}(\nabla^Q)$ can be computing using [Qu], [BGS].

Let $\nabla^{\mathcal{L}}$ denote the Chern connection for $(\mathcal{L}\,,h)$. Its curvature will be denoted by $\mathcal{K}(\nabla^{\mathcal{L}})$.

Proposition 2.2. *The two $(1,1)$–forms $\mathcal{K}(\nabla^Q)$ and $\mathcal{K}(\nabla^{\mathcal{L}})$ on M coincide.*

Proof. The Chern connection on the holomorphic Hermitian line bundle $(L_0\,,h_0)$ (respectively, $(L\,,H)$) will be denoted by ∇^{L_0} (respectively, ∇^L). Let $\mathcal{K}(\nabla^{L_0})$ (respectively, $\mathcal{K}(\nabla^L)$) be the curvature of ∇^{L_0} (respectively, ∇^L). From the definition of H (see (2.3)) it follows immediately that

$$\mathcal{K}(\nabla^L)\,=\,p_1^*\mathcal{K}(\nabla^{L_0})+p_2^*\mathcal{K}(\nabla^{\mathcal{L}})\,. \tag{2.6}$$

Let $\mathcal{K}(\omega_X) \in C^\infty(X; \Omega_X^{1,1})$ be the curvature of TX for the Kähler form ω_X.

A theorem due to Quillen and Bismut–Gilet–Soulé says that $\mathcal{K}(\nabla^Q)$ is given by the following fiber integral along X

$$\mathcal{K}(\nabla^Q) \,=\, -\frac{1}{2\pi\sqrt{-1}}\left(\int_{(X\times M)/M} (\mathcal{K}(\nabla^L) + \frac{1}{2}\mathcal{K}(\nabla^L)^2) \wedge (1 + \frac{1}{2}p_1^*\mathcal{K}(\omega_X))\right)_2 \tag{2.7}$$

[BGS, p. 51, Theorem 0.1], [Qu], where $(\beta)_2$ denote the component of the differential form β of degree two; note that $\frac{1}{4\pi\sqrt{-1}}\mathcal{K}(\omega_X)$ is the Todd form on X for the Kähler form ω_X that represents the Todd class $\frac{1}{2}c_1(TX)$. Using (2.6), the expression in (2.7) reduces to

$$2\pi\sqrt{-1}\cdot\mathcal{K}(\nabla^Q) \,=\, -\mathcal{K}(\nabla^{\mathcal{L}})\cdot\int_X (\mathcal{K}(\nabla^{L_0}) + \frac{1}{2}\mathcal{K}(\omega_X))\,. \tag{2.8}$$

Now note that

$$\frac{1}{2\pi\sqrt{-1}}\int_X (\mathcal{K}(\nabla^{L_0}) + \frac{1}{2}\mathcal{K}(\omega_X)) \,=\, \text{degree}(L_0) + \frac{1}{2}\text{degree}(TX) \,=\, g-2+1-g \,=\, -1\,.$$

Using this, from (2.8) we conclude that $\mathcal{K}(\nabla^Q) \,=\, \mathcal{K}(\nabla^{\mathcal{L}})$. □

The Hermitian structure L_0 and the Kähler form ω_X together produce an inner product on $H^1(X,\, L_0)$. This inner product defines a Hermitian structure h_ξ on the holomorphic line bundle ξ in (2.4). Note that the Chern connection on ξ for h_ξ is flat.

The Hermitian structure h on $\mathcal{L}$ and the Hermitian structure h_ξ on ξ together produce a Hermitian structure $\widetilde{h}$ on $\mathcal{L}\otimes\xi$.

Proposition 2.3. *For the isomorphism τ in* (2.5)*, there is a positive real number t such that $\tau^*\widetilde{h} \,=\, t\cdot h_Q$.*

Proof. There is a real valued C^∞ function f on M such that

$$\tau^*\widetilde{h} \,=\, \exp(f)\cdot h_Q\,.$$

From Proposition 2.2 it follows the two holomorphic Hermitian line bundles $(\mathcal{Q}\,, h_Q)$ and $(\mathcal{Q}\,, \exp(f)\cdot h_Q)$ have the same curvature. This implies that f is a harmonic function. Since M is compact and connected, any harmonic function on it is a constant one. □

Theorem 2.4. *The two G–equivariant holomorphic Hermitian line bundles $(\mathcal{L}\,, h)$ and $(\mathcal{Q}\,, h_Q)$ are isomorphic.*

Proof. Consider the isomorphism

$$\frac{1}{\sqrt{t}}\cdot\tau \,:\, \mathcal{Q} \,\longrightarrow\, \mathcal{L}\otimes\xi\,,$$

where τ is the isomorphism in (2.5), and t is the constant in Proposition 2.3. From Lemma 2.1 and Proposition 2.3 it follows immediately that this is a G–equivariant holomorphic isomorphism that takes the Hermitian structure h_Q on $\mathcal{Q}$ to the Hermitian structure $\widetilde{h}$ on $\mathcal{L}\otimes\xi$.

The two G–equivariant holomorphic Hermitian line bundles $(\mathcal{L}\,, h)$ and $(\mathcal{L}\otimes\xi\,, \widetilde{h})$ are clearly isomorphic. Therefore, the two G–equivariant holomorphic Hermitian line bundles $(\mathcal{L}\,, h)$ and $(\mathcal{Q}\,, h_Q)$ are isomorphic. □

Acknowledgements

The author is grateful to the Harish–Chandra Research Institute for hospitality while the work was carried out. The author acknowledges the support of the J. C. Bose Fellowship.

References

[BGS] J.-M. Bismut, H. Gillet and C. Soulé, Analytic torsion and holomorphic determinant bundles. I. Bott-Chern forms and analytic torsion, *Comm. Math. Phys.* **115** (1988), 49–78.

[DM] R. Dey and V. Mathai, Holomorphic Quillen determinant line bundles on integral compact Kähler manifolds, *Quat. Jour. Math.* **64** (2013), 785–794.

[Ha] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977.

[Qu] D. G. Quillen, Determinants of Cauchy-Riemann operators on Riemann surfaces, *Funct. Anal. Appl.* **19** (1985), 37–41.

School of Mathematics, Tata Institute of Fundamental Research, Homi Bhabha Road, Bombay 400005, India

E-mail address: indranil@math.tifr.res.in